

Q D

305

A2Ms

UC-NRLF



B 2 736 694

YD 04968

EXCHANGE



EX LIBRIS

529
1454

TC
[Reprinted from the Journal of the American Chemical Society,
Vol. XLIII, No. 12, December, 1921]

Experimental Studies on Cystine

DISSERTATION *cat. 192*

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIRE-
MENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
IN THE FACULTY OF PURE SCIENCE IN
COLUMBIA UNIVERSITY

BY

ALICE RANNEY THOMPSON MERRILL, B.S., A.M.

New York City

1921



The Jackson Press, Kingston
1921

[Reprinted from the Journal of the American Chemical Society,
Vol. XLIII, No. 12, December, 1921]

Experimental Studies on Cystine

DISSERTATION

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIRE-
MENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
IN THE FACULTY OF PURE SCIENCE IN
COLUMBIA UNIVERSITY

BY

ALICE RANNEY THOMPSON MERRILL, B.S., A.M.

New York City

1921

The Jackson Press, Kingston
1921

70 1000
A000000000

QJ 305
A2 MS

ACKNOWLEDGMENTS

The author wishes to express her appreciation to Professor H. C. Sherman, who suggested this investigation.

To Professors H. C. Sherman, A. W. Thomas, and H. T. Beans, and others is due much gratitude for helpful advice and suggestions throughout the work.

EXCHANGE

[Reprinted from the Journal of the American Chemical Society,
Vol. XLIII. No. 12. December, 1921.]

[CONTRIBUTION FROM THE CHEMISTRY DEPARTMENT OF COLUMBIA
UNIVERSITY, No. 380.]

EXPERIMENTAL STUDIES ON CYSTINE

BY ALICE R. THOMPSON MERRILL.

Received August 29, 1921.

In preparing cystine for biological experiments it is of importance to know the conditions which favor the precipitation of pure cystine.

It has been shown by Mörner,¹ Embden,² Abderhalden³ and others that cystine may be obtained by the acid hydrolysis of horn, wool and other protein substances containing cystine in combination. When the acid solution is neutralized with alkali or brought to weak acidity with acetic acid after neutralization, the cystine precipitates on standing.

Folin⁴ adds sodium acetate to the acid solution of cystine using congo red as indicator. The sodium acetate acts as a buffer salt. He states that this method gives satisfactory results but does not give quantitative directions for the method.

The purpose of this investigation was (1) to determine the concentration of hydrogen ion of solutions of cystine which insures maximum precipitation of cystine (2) to determine the amounts of sodium acetate to add to the hydrolyzed solution of wool in order to obtain the best yields of cystine and (3) to secure data in regard to purification of the crude cystine from the hydrolyzed solutions. The yield of cystine obtained by Folin's method is compared to the yield obtained by earlier methods.

Experimental Procedure

The wool used in this investigation was analysed and found to contain 3.1% sulfur and 11.6% moisture. Two portions of 750 g.

¹Mörner, Z. physiol. Chem., 28, 595 (1899).

²Embden, *ibid.*, 32, 94 (1900).

³Abderhalden, *ibid.*, 52, 348 (1907).

⁴Folin, J. Biol. Chem., 8, 9 (1910).

each were boiled with 1500 cc. of conc. hydrochloric acid in separate 3-liter flasks with large necks fitted with reflux water condensers. The solutions in the flasks were heated in oil-baths at a temperature of about 115°. In spite of the condensers, hydrogen chloride fumes poured out through the water-cooled tubes during the first part of the hydrolysis and continued to do so until the acid mixture approached the concentration of the acid-water mixture of constant boiling point. The time of hydrolysis was counted from the time at which all the wool was pushed into the hot acid. The hydrolysis was run for 3 hours, when the biuret test became negative.

The solutions on cooling and thorough mixing had a volume of 4065 cc. and an acidity of 6.9 molar strengths. Aliquots of 500 cc. representing 188.0 g. of wool were taken and sodium acetate trihydrate crystals added to each aliquot as shown in Table I. The solutions were heated to boiling and boiled for 5 minutes with constant stirring. After standing for 3 days the precipitate of crude cystine was filtered by suction, removed, rubbed up with cold water, filtered and washed once on a Büchner funnel. This process was repeated. The undiluted mother liquors were kept in glass-stoppered bottles and the washings kept separately.

The mother liquors on standing for two weeks were filtered by suction and the second precipitates washed and filtered as the first were.

The precipitates were air dried, ground to a powder and kept in desiccators until they came to constant weights.

Sulfur was determined in these weighed precipitates by Osborne's peroxide method. Sulfur was also determined in pure cystine by the Carius method and the results compared with those obtained by the fusion method. It was found that the Osborne method gave results from about 0.2 to 0.3% lower than those obtained by the Carius method on cystine containing 26.69% sulfur. Half-gram samples were fused and an aliquot representing 0.1 g. taken for the precipitation of barium sulfate. Great care was taken to determine and subtract the sulfur in reagents used, to insure barium sulfate precipitates from reduction during ignition and to make determinations in duplicate.

In Table I are shown the grams of sulfur in each precipitate of crude cystine as calculated from the analysis; also the percentages of cystine obtained from the wool. It is considered here that the

sulfur determined represents cystine sulfur. This assumption seems correct since these precipitates were almost completely soluble in 1.8 M (6.5%) hydrochloric acid and fairly insoluble in cold water as is cystine also, and the yield of pure cystine obtained by experiment corresponds closely to the yield indicated by calculations from sulfur as determined in the crude precipitates.

TABLE I

Sulfur in Crude Cystine Precipitates from Aliquots Representing 188 g. of Wool

NaC ₂ H ₃ O ₂ (3H ₂ O)	(1) S in 1st ppt.	(2) S in 2nd ppt.	Total S in 1 and 2.	Total sulfur in wool.	Total cystine in wool calc. from sulfur found
G.	G.	G.	G.	%.	%.
350	0.077	0.615	0.692	0.368	1.379
450	0.648	0.975	1.623	0.865	3.241
500	1.273	0.746	2.019	1.075	4.028
550	1.260	0.705	1.965	1.046	3.919
650	2.048	0.112	2.160	1.150	4.309
750	2.268	0.055	2.323	1.236	4.631
850	1.706	0.093	1.799	0.957	3.586
950	1.976	0.151	2.127	1.132	4.242

It is evident from Table I that 350 g. of sodium acetate trihydrate precipitated only traces of cystine after standing for 3 days, while 550 g. of sodium acetate gave a yield of only a little over one-half of the cystine given by 650 g. of acetate; 750 g. of the salt gave the maximum yield of cystine.

The sulfur in the precipitates obtained from the mother liquors which stood for 2 weeks after the first filtration was greatest in those in which insufficient sodium acetate had been added to secure good yields of cystine in the first precipitation.

The total yield of cystine in the first and second precipitates from the aliquot in which 500 g. of acetate had been added was almost as large as with greater amounts of sodium acetate, but required 2 weeks' standing instead of 3 days.

Determination of Hydrogen-ion Concentrations

In order to determine the concentration of the hydrogen ion more accurately in the cystine solutions, a set of buffer solutions was made up. The acetic acid-sodium acetate mixtures of Walpole⁵

⁵Walpole, J. Chem. Soc., 105, 2501 (1914).

were used in place of the potassium hydrogen phthalate solutions recommended by Clark and Lubs,⁶ as the latter gave a decided drift when determining the hydrogen-ion concentration by the electrometric method, due possibly to reduction occurring. The set of standard solutions of Clark and Lubs⁶ was also made up and used with the one exception stated above. The hydrogen-ion concentrations of these standard buffer solutions were determined directly in the solutions using the hydrogen electrode. By use of the indicators listed by Clark and Lubs the standard solutions were compared colorimetrically with the cystine solutions. It was considered inadvisable to determine the hydrogen-ion concentration in the cystine solutions by the electrometric method because of probable reduction by the hydrogen electrodes.

Since the hydrolyzed solutions were colored dark brown they were diluted to 5 times the original volume as recommended by Walpole⁵. Measured amounts of indicator were added to given volumes of cystine solutions and compared with standard solutions containing the same amount of indicator. That the tints might be comparable the standard solutions plus the indicator were superimposed against the dark solution according to the method of Walpole⁷ but modified as follows.

Modified Comparator Used.—Difficulty was experienced in the use of the comparator of Humitz, Mayer and Ostenberg⁸ due to the reflection of the light from the round sides of the glass tubes. It was believed that glass cells with flat sides and fitted together in pairs would give better results. Four flat-sided glass cells generally used with colorimeters were therefore picked out. These cells were 10×35×53 mm. inner dimensions and were placed together in pairs, the broad sides fitted and held together by elastic bands. The cells were filled with solutions and superimposed according to the Walpole combination.

The cells were placed on dull black cardboard, placed before the screen described by Clark and Lubs⁹ and lighted by 4 electric bulbs. A dull black cardboard with vertical rectangular windows cut close together was drawn over the cells and the color of each as seen through the apertures compared. The cells were also com-

⁶Clark and Lubs, J. Biol. Chem., 25, 479 (1916).

⁷Walpole, Biochem. J., 5, 207 (1910).

⁸Humitz, Meyer and Ostenberg, Proc. Soc. Exp. Biol. Med., 13, 24 1915.

⁹Clark and Lubs, J. Bact., 2, 134 (1917).

pared in daylight when suitable indicators were used. This substitute for the black box comparator was very satisfactory as well as simple to use.

Concentration of Hydrogen Ion of Hydrolyzed Solutions.—Using the modified comparator and brom phenol blue indicator, it was found that the hydrolyzed solution giving maximum yields of cystine after addition of sodium acetate showed a concentration of hydrogen ion equal to 10^{-4} .

It was noticed that congo red gives a red color when sodium acetate is added to a hydrolyzed solution in amounts still insufficient to give maximum yields of cystine, and therefore this red color does not indicate that sufficient sodium acetate has been added.

Hydrolysis of Wool

A series of hydrolyses of wool was next made to determine the effect of increasing the length of time of hydrolysis on the yield of cystine. Portions of 500 g. of wool were boiled with 1000 cc. of conc. hydrochloric acid for different periods of time and 750 g. of sodium acetate added to aliquots representing 185 g. of wool. Sulfur was determined in the precipitates formed on standing 3 days and also on precipitates in the mother liquor. Alcohol was also added to the mother liquors until they became turbid, and the precipitates thus formed were analysed.

TABLE II

Hydrolysis with Concentrated Hydrochloric Acid. Sulfur and Cystine in Precipitates from Aliquots Representing 185 g. of Wool

Time of hydrolysis.	Sulfur.	Calc. cystine in wool.
Hours	G	%
3	1.618	3.27
4	2.037	4.12
5	2.120	4.29
6	2.135	4.32
12	2.578	5.22
74	0.120	0.24

A study of Table II suggests that 3 hours' hydrolysis is not a sufficient length of time but that there is an increased yield of cystine up to the 12-hour period. The 74-hour period gives practically no yield of cystine.

The addition of alcohol gave a slight increase in yield.

Hydrolysis was carried out with 6.8 *M* (22%) hydrochloric acid. Two hundred and fifty g. of wool was boiled with 900 cc. of dil. acid for 12 hours. From an aliquot representing 185 g. of wool only 1.532 g. of sulfur was precipitated, but on adding alcohol until the solution became turbid 1.354 g. of sulfur was further precipitated, giving a total calculated yield of cystine equal to 5.84% of the wool. This indicates that hydrolysis with dilute hydrochloric acid gives good yields of cystine, but as the mother liquors obtained from dilute acid contain less hydrolyzed wool per volume they tend to retain more cystine in solution after neutralization. The advantage of using dil. hydrochloric acid is that while pushing the wool into the hot acid no strong hydrogen chloride fumes are evolved to burn the hands.

Preparation of Pure Cystine

The quantitative yields of pure cystine obtained from the crude cystine precipitated by sodium acetate as above were determined.

Two portions of 500 g. of wool were boiled with 1000 cc. of conc. hydrochloric acid each, for 6 hours as usual. Sodium acetate sufficient to give a maximum yield of cystine was added. Each washed and dried precipitate was treated as follows. The precipitate was dissolved in 250 cc. of 1.8 *M* (6.5%) hydrochloric acid, filtered by suction, the insoluble portion digested with 2 portions of 50 cc. of hydrochloric acid and filtered. Norite freed from iron and calcium was added equal to 50 g. on the water-free basis, to the combined filtrates, heated and filtered. The Norite was boiled with 2 portions of 100 cc. of 5% hydrochloric acid¹⁰ and filtered. The filtrates were practically colorless. Five hundred cc. of hot sodium acetate trihydrate solution (1:1) was added to the hot combined acid solution. Almost pure cystine was precipitated. Analysis of the white crystals of cystine showed them to be 98.6% pure. By analysis for sulfur before and after use it was found that the Norite had adsorbed 0.313 g. of sulfur.

The yield of cystine thus obtained was 4.3% of the wool and practically equal to the yield as calculated from sulfur determined in the crude precipitate.

Analysis of Pure Cystine

Pure cystine was obtained by repeatedly dissolving in 1.8 *M* (6.5%) hydrochloric acid and precipitating with sodium acetate solution. Sulfur in one sample was found by 3 determinations by the Carius method to be 26.66%, 26.63% and 26.86% respectively; the calculated sulfur is 26.69%. By the peroxide method with the same sample 26.37% was obtained.

¹⁰Miss Dennis recommends that bone-black used in decolorizing these precipitates be boiled with dilute acid. *J. Biol. Chem.*, 9, 369 (1911).

Cystine Obtained by the Method used by Abderhalden

Two hundred and fifty g. of wool was boiled under a reflux condenser with 900 cc. of 6.8 *M* (22%) hydrochloric acid for 12 hours, and evaporated to a gummy mass *in vacuo*. This was dissolved in water, made up to 1000 cc., decolorized with Norite, and 100 g. of sodium hydroxide in solution added to the cold filtrate. After standing for 3 days the fine white crystals were filtered with suction. The concentration of hydrogen ion in the filtrate was found to be $10^{-4.4}$. The filtrate was evaporated *in vacuo* and a further yield of cystine obtained.

TABLE III

Cystine by Abderhalden's Method

	Sulfur. G.	Cystine. G.	Cystine in Wool. %
1st Ppt.	2.910	10.904	4.36
2nd Ppt.	1.429	5.354	2.14
Total	4.339	16.258	6.50

The total yield of cystine obtained by this method was 6.50% of wool. Here again it is noticed that concentration of the mother liquor results in further precipitation of cystine from solution.

The advantage of using sodium hydroxide solution over that of using sodium acetate is that less salt is formed in the mother liquor and concentration of the latter is easier. In Folin's method the mother liquors are saturated with salt.

The Determination of the Concentrations of Hydrogen Ion in Solutions in which Maximum Yields of Cystine are Obtained

The hydrogen-ion concentration of the solutions of hydrolyzed wool which on addition of sodium acetate had given the maximum yield was found to be 10^{-4} .

Of special interest are the results obtained in determining the concentration of hydrogen ion in solutions which give maximum and minimum precipitates of cystine from a solution of pure cystine. Three series were run to obtain these results. In each, 10 g. of pure cystine was dissolved in 1.8 *M* hydrochloric acid and made up to 500 cc. with 1.8 *M* hydrochloric acid.

Fifty cc. aliquot portions containing 1 g. of cystine each were run into separate beakers. Sodium acetate trihydrate in solution (1:1) was added from a buret in various amounts to each beaker and distilled water added to make the volume in each case 150 cc. In another series 4.34 *M* sodium hydroxide solution was added first to 50 cc. of the acetate

solution and rinsed with water into the aliquots of cystine solution; the volume was then made up as before to 150 cc. After standing for 48 hours at room temperatures the precipitates formed were filtered into weighed Gooch crucibles with asbestos pads, washed thrice with cold water and once each with alcohol and ether. The precipitates were dried at 100°, cooled in the desiccator and weighed.

The concentration of hydrogen ion was determined in the filtrates kept separate from the washings. Standard buffer solutions, the hydrogen-ion concentrations of which had been determined by the electrometric method, were used for comparison in determining the concentration of hydrogen ion in the filtrates by the colorimetric method.

In Table IV are shown the amounts of reagents used, the percentages of cystine obtained and the corresponding concentrations of hydrogen ion.

The maximum yield of cystine was 96.4%. It is clear that addition of the 1:1 sodium acetate solution to 1.8 M hydrochloric

TABLE IV

Percentages of Total Cystine in Solution Precipitated and Concentrations of Hydrogen Ion

NaC ₂ H ₃ O ₂ ·3H ₂ O sol. 1:1.	NaOH 4.34 M	Cystine precipitated.	Log C _H ⁺ .
Cc.	Cc.	%	
10.0	...	00.00
12.0	...	00.00
14.0	...	00.00	—1.0
16.0	...	11.93	—1.1
16.5	...	16.06	—1.2
17.0	...	53.11	—1.3
17.5	...	62.18	—1.4
18.0	...	75.85	—1.6
19.0	...	93.97	—2.3
20.0	...	96.22	—3.4
30.0	...	96.37	—4.3
40.0	...	96.10	—4.5
50	...	96.29	—4.6
50	18.0	94.54	—6.9
50	18.5	84.91	—8.3
50	19.0	54.64	—8.9
50	19.5	30.77	—8.9
50	20.0	2.67	—9
50	20.5	0	—9




PLATE I

Fig. 1

acid gives a maximum yield when 20 to 30 cc. of the acetate solution is added to 50 cc. of the dil. hydrochloric acid solution of cystine. As the iso-electric point of tyrosine, calculated from the values $ka=4\times 10^{-9}$ and $kb=2.6\times 10^{-12}$ as determined by Kanitz¹¹ and using the formula¹²

$$L = \sqrt{\frac{ka}{kb}} \times kw$$

is $10^{-5.4}$, it is well to precipitate the cystine at a hydrogen-ion concentration greater than that of $10^{-5.4}$ to obtain it free from tyrosine.

The curve in Fig. 1 follows the values of Table IV. The iso-electric zone for cystine as indicated is a broad one and shows that cystine can be precipitated in solutions of quite varying strengths.

Summary

It has been shown that the zone of hydrogen-ion concentration most favorable for the precipitation of cystine from solution is be-

¹¹Kanitz, Z. physik. Chem., 47, 476 (1906).

¹²Michaelis and Mostynski, Biochem. Z., 24, 79-91 (1910).

Michaelis and Davidsohn, *ibid.*, 30, 143-50 (1911).

tween 10^{-3} and 10^{-6} . To obtain cystine free from tyrosine it is advisable to precipitate it from solution at a concentration of hydrogen ion of about 10^{-3} .

The quantity of sodium acetate required as a buffer in the hydrolyzed solution of wool was determined. When 100 g. of wool is hydrolyzed with 200 cc. of conc. hydrochloric acid, sp. gr. 1.19, 750 g. of sodium acetate trihydrate crystals should be added to 500 cc. of the hydrolyzed solution to insure maximum precipitation of cystine. The concentration of hydrogen ion was determined and found to be 10^{-4} . The highest yield of cystine thus obtained was 5.2% of the weight of the wool taken.

When sodium hydroxide solution was added to a hydrolyzed solution which had been evaporated *in vacuo* and taken up in water, the yield of cystine obtained was 6.5% of the wool. The filtrate contained less salt than that obtained by Folin's method and could therefore be concentrated with less difficulty. Cystine was obtained pure from the crude cystine preprecipitated by sodium acetate from the hydrolyzed solution of wool and the percentage yield of pure cystine actually obtained agreed with the yield calculated by determination of sulfur in the crude precipitate.

Comparisons were made between the yields of cystine obtained after boiling wool with conc. hydrochloric acid, sp. gr. 1.19, for from 3 to 12 hours with the yield obtained using 6.8 *M* (22%) hydrochloric acid for hydrolysis. It was found that boiling wool with conc. hydrochloric acid for 12 hours gave the highest yield of cystine. When about 20% hydrochloric acid was used, the solution was so dilute that alcohol had to be added to the buffered mixture. When alcohol was added, the yield of cystine was as high from the dilute acid hydrolysis as with the concentrated. Dil. hydrochloric acid does not evolve the strong acid fumes that the concentrated acid does and is therefore more convenient to handle although it takes more space.

The amount of cystine adsorbed by the Norite used in decolorizing the crude cystine was found to be small when the Norite had been boiled with dil. hydrochloric acid and filtered after decolorizing.

In determining the hydrogen concentration of dark colored solutions a modification was made in the comparator used by Humitz, Meyer and Ostenberg.

NEW YORK CITY.

VITA

Alice R. Thompson Merrill was born in Hoboken, N.J., December 22, 1883. She graduated from the College of Agriculture, University of California, in January, 1906.

She was assistant soil chemist in the College of Agriculture, University of California, until June, 1906, and then accepted a position as assistant chemist in the U. S. Agricultural Experiment Station in Hawaii. In her work at this Station she made a number of investigations in regard to Hawaiian soil, plants, and food, and published several articles in the Station Reports and Bulletins and also in Chemical Journals.

In 1910-11 she spent a year at Columbia University studying under the Faculty of Pure Science and obtained the degree of M.A. On her return to Hawaii, she renewed her work at the U. S. Agricultural Station there and published several bulletins and articles.

In 1916 she accepted a position in the Bureau of Chemistry, Department of Agriculture, Washington, D.C., and remained there until 1917, when she returned to graduate study in chemistry under the Faculty of Pure Science, Columbia University.

The titles of her publications are as follows:

Hawaiian Honeys. Hawaii Agri. Exp. Sta. Bull. 17, pt. II, 1908.

Report of Assistant Chemist, *ibid.*, 1907.

Report of Assistant Chemist, *ibid.*, 1908.

The Determination of Sulphur and Chlorine in the Rice Plant, Annual Report Hawaii Agri. Exp. Sta., 1912, also in J. Amer. Chem. Soc. 35, 1628 (1913).

The Extraction and Use of Kukui Oil. Hawaii Agri. Exp. Sta., Press Bull. 39, pt. II, 1912, also in J. Ind. Eng. Chem. 5, 644, 1913.

Organic Phosphoric Acid of Rice, U.S.D.A., J. Agri. Research, 3, 425.

Composition of Hawaiian Fruits and Nuts. Annual Report Hawaii Agri. Exp. Sta., 62, 1914.

Chemical Studies of the Efficiency of Legumes as Green Manures in Hawaii. Hawaii Agri. Sta. Bull. 43.

**THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW**

**AN INITIAL FINE OF 25 CENTS
WILL BE ASSESSED FOR FAILURE TO RETURN
THIS BOOK ON THE DATE DUE. THE PENALTY
WILL INCREASE TO 50 CENTS ON THE FOURTH
DAY AND TO \$1.00 ON THE SEVENTH DAY
OVERDUE.**

MAR 24 1933

LD 21-50m-1,'33

Photomount
Pamphlet
Binder
Gaylord Bros.
Makers
Syracuse, N. Y.
PAT. JAN 21, 1908

YD 04968

507775

QD305

A2M5

UNIVERSITY OF CALIFORNIA LIBRARY

